Dynamics of Dirac solitons in networks

K.K. Sabirov[a,b], D.B. Babajanov[a], D.U. Matrasulov[a] and P.G. Kevrekidis[c1]

[a] *Turin Polytechnic University in Tashkent, 17 Niyazov Str., 100095, Tashkent, Uzbekistan*

[b] *Tashkent University of Information Technologies, 102 Amir Temur Str., 100200, Tashkent, Uzbekistan*

[c] *Department of Mathematics and Statistics, University of Massachusetts, Amherst, Massachusetts 01003-4515 USA*

We study dynamics of Dirac solitons in prototypical networks modeling them by the nonlinear Dirac equation on metric graphs. Soliton solutions of the nonlinear Dirac equation on simple metric graphs are obtained. It is shown that these solutions provide reflectionless vertex transmission of the Dirac solitons under suitable conditions. The constraints for bond nonlinearity coefficients, allowing reflectionless transmission over a Y-junction are derived. The analytical results are confirmed by direct numerical simulations.

arXiv:1701.05707v1 [nlin.PS] 20 Jan 2017

I. INTRODUCTION

Nonlinear evolution equations on networks have attracted much attention recently[1–10]. Such interest is caused by a broad variety of potential applications of the nonlinear wave dynamics and soliton transport in networks, such as Bose-Einstein condensates (BECs) in branched traps, Josephson junction networks, DNA double helix, polymer chains, etc.

Despite the rapidly growing interest in wave dynamics on networks, most of the studies are mainly focused on nonrelativistic wave equations such as the nonlinear Schrödinger equation[1–7,10]. Nevertheless, there is a number of works on the sine-Gordon equation in branched systems[8,9,11]. However, relativistic wave equations such as the nonlinear Klein-Gordon and Dirac equations are important in field theory and condensed matter physics and hence exploring them on metric graphs is of interest in its own right. The latter consist of a system of bonds which are connected at one or more vertices (branching points). The connection rule is associated with the topology of a graph. When the bonds can be assigned a length, the graph is called a metric graph.

In this paper we address the problem of the nonlinear Dirac equation on simple metric graphs by focusing on conservation laws and soliton transmission at the graph vertices. Our prototypical example will be the Y-junction. Early studies of the nonlinear Dirac equation date back to Thirring[12] and Gross-Neveu[13] models of field theory. Integrability, nonrelativistic limit and exact solutions of the relevant models have been considered in[14–16]. A potential application of the nonlinear Dirac equation to neutrino oscillations was discussed in[17]. Recently, the possibility of experimental realization of Dirac solitons in Bose-Einstein condensates in honeycomb optical lattices was discussed in[19] where the nonlinear Dirac equation was derived. Further analysis of this setting in Refs.[20–22,24] has excited a rapidly growing interest in the nonlinear Dirac equation and its soliton solutions (see, e.g.[25]-[47]). In[25,26,29–32], a detailed study of soliton solutions for different types of nonlinearity, their stability and discussions of conservation laws are presented. These developments have, in turn, had an impact also on the mathematical literature where the stability of solutions for different forms of the nonlinear Dirac equation was explored in[27,28,34]. In[35]-[39], the stability of Gross-Neveu solitons in both 1d and 3d and for other nonlinear Dirac models an important energy-based stability criterion was developed. The numerical corroboration of the stability for solitary waves and vortices in 2D nonlinear Dirac models of the Gross-Neveu type was considered quite recently in[33].

Dirac solitons in branched systems can, in principle, be experimentally realized in different systems of optics, but also envisioned elsewhere (e.g. in atomic physics etc.). A relevant such possibility consists of the discrete waveguide arrays of[42,47] that can be formulated as a branched system to be described (in the appropriate long wavelength limit[42])

by a nonlinear Dirac equation on metric graphs. Such networks in optical systems have been proposed, e.g., earlier in[48]. More exotic possibilities can be envisioned in branched networks of honeycomb (i.e., armchair nanoribbon) optical lattices for atomic BECs[21], although we will not focus on the latter here.



FIG. 1. Metric star graph

In the present work, we focus on a prototypical example of exact solutions and transport of Dirac solitons through the network vertices. In particular, we show that under certain constraints, exact soliton solutions of the nonlinear Dirac equation on simple metric graphs can be obtained. The identified soliton solutions provide reflectionless transmission of the Dirac solitons at the vertices. This renders possible the tuning of the transport properties of a network in such a way that it can provide ballistic transport of the Dirac solitons. This paper is organized as follows: In the next section we give the formulation of the problem on a metric star graph which includes the derivation of the vertex boundary conditions. Section III presents soliton solutions of the nonlinear Dirac equation on a metric star graph for fixed and moving frames. Also, an analysis of the vertex transmission of the Dirac solitons on the basis of the numerical solution of the nonlinear Dirac equation is presented. In section IV we extend the treatment to a metric tree graph and discuss the extension to other simple topologies. Finally, section V presents some concluding remarks.

II. CONSERVATION LAWS AND VERTEX BOUNDARY CONDITIONS

The nonlinear Dirac system that we are going to explore, i.e., specifically the Gross-Neveu model, follows from the Lagrangian (in the units $\hbar = c = 1$)[25,26,29]

$$L = \frac{i}{2}\bar{\Psi}\gamma^{\mu}\partial_{\mu}\Psi - m\bar{\Psi}\Psi + \frac{g^2}{2}\left(\bar{\Psi}\Psi\right)^2, \quad \mu = 0, 1, \tag{1}$$

and describes interacting Dirac fields. Field equations for this Lagrangian lead to the nonlinear Dirac model of the form

$$\left(i\gamma^{\mu}\partial_{\mu} - m\right)\Psi + g^2\left(\bar{\Psi}\Psi\right)\Psi = 0, \tag{2}$$

where

$$\Psi(x,t) = \begin{pmatrix} \phi(x,t) \\ \chi(x,t) \end{pmatrix}; \quad \bar{\Psi} = \Psi^{\dagger}\gamma^0, \tag{3}$$

and $\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, $\gamma^1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

For a metric star graph consisting of three semi-infinite bonds (see, e.g., Fig. 1), the Lagrangian density of the nonlinear Dirac field for each bond can be written as L_j where j parametrizes the bond and L for each is of the form of Eq. (1). The field equation following from this Lagrangian density can be written as Eq. (2) before, where the spatial coordinates are defined as $x_1 \in (-\infty, 0]$ and $x_{2,3} \in [0, \infty)$, while 0 coincides with the graph vertex.

Formulation of an evolution set of equations on metric graphs requires imposing vertex boundary conditions which provide "gluing" of the graph bonds at the graph vertices. For the linear Dirac equation on metric graph studied earlier by Harrison and Bolte in[49], the vertex boundary conditions have been derived from the self-adjointness of the Dirac operator on a graph. In that case such conditions led to Kirchhoff rules and continuity of the wave function at the graph vertex[49]. However, for nonlinear problem in order to derive vertex BCs, it is arguably more natural to use suitable conservation laws of the nonlinear flow, which give rise to appropriate BCs. For the nonlinear Dirac system fundamental conservation laws can be presented in terms of the momentum-energy tensor given by[25,26]

$$T^j_{\mu\nu} = \frac{i}{2}[\bar{\Psi}_j\gamma^\mu\partial^\nu\Psi_j - \partial^\nu\bar{\Psi}_j\gamma^\mu\Psi_j] - g^{\mu\nu}L_j,$$

where $g^{\mu\nu} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

The energy on each bond of a star graph can be written as

$$E_j = -\int_{b_j} T^j_{00}dx = -\int_{b_j}[\frac{i}{2}[\bar{\Psi}_j\gamma^1\partial_x\Psi_j - \partial_x\bar{\Psi}_j\gamma^1\Psi_j] - m\bar{\Psi}_j\Psi_j + \frac{g_j^2}{2}(\bar{\Psi}_j\Psi_j)^2]dx, \tag{4}$$

where integration is performed along each bond. Here, to derive the vertex boundary conditions (VBC) we use conservations of particle current and energy. The total particle current for the star graph is defined as

$$J = J_1 + J_2 + J_3,$$

where the current for each bond is given by

$$J_j = \int_{b_j} \left(|\phi_j|^2 + |\chi_j|^2\right) dx. \tag{5}$$

The current conservation, $\dot{J} = 0$ yields the following vertex boundary condition:

$$\mathrm{Re}\,[\phi_1\chi_1^*]|_{x=0} = \mathrm{Re}\,[\phi_2\chi_2^*]|_{x=0} + \mathrm{Re}\,[\phi_3\chi_3^*]|_{x=0}\,. \tag{6}$$

Here we used the asymptotic conditions

$$\Psi_1 \to 0 \text{ at } x \to -\infty \text{ and } \Psi_{2,3} \to 0 \text{ at } x \to \infty. \tag{7}$$

The energy conservation, $\dot{E} = 0$ leads to

$$\mathrm{Im}\,[\phi_1\partial_t\chi_1^* + \chi_1\partial_t\phi_1^*]|_{x=0} = \mathrm{Im}\,[\phi_2\partial_t\chi_2^* + \chi_2\partial_t\phi_2^*]|_{x=0} + \mathrm{Im}\,[\phi_3\partial_t\chi_3^* + \chi_3\partial_t\phi_3^*]|_{x=0}\,. \tag{8}$$

As per the above analysis, the VBC for NLDE on metric graphs have been obtained from the energy and current conservation. However, the VBC given by Eqs. (6) can be fulfilled, if the following linear relations at the vertices are imposed (see Appendix for details):

$$\begin{aligned} \alpha_1\phi_1|_{x=0} &= \alpha_2\phi_2|_{x=0} + \alpha_3\phi_3|_{x=0}, \\ \frac{1}{\alpha_1}\chi_1|_{x=0} &= \frac{1}{\alpha_2}\chi_2|_{x=0} = \frac{1}{\alpha_3}\chi_3|_{x=0}, \end{aligned} \tag{9}$$

where $\alpha_1, \alpha_2, \alpha_3$ are the real constants which will be determined below. In the following we will use Eqs. (9) as the vertex boundary conditions for Eq. (2) on a metric star graph. When applied to the exact soliton solutions of the NLDE, the VBC will lead to algebraic conditions connecting α_j and g_j, as will be derived in the next section.

III. SOLITON DYNAMICS AND VERTEX TRANSMISSION

In[25,26,29] the soliton solutions of the nonlinear Dirac equation have been obtained in the form of standing wave solutions asymptoting to zero as $x \to \pm\infty$. Here, we use the same definition for soliton solutions of the nonlinear Dirac equation on the metric star graph presented in Fig. 1. We note in passing that for soliton solutions of the NLDE derived in the Refs.[25,26,29] all elements of the energy-momentum tensor, except T_{00} become zero, i.e., $T_{01} = T_{10} = T_{11} = 0$.

We look for the soliton solution of NLDE on a metric star graph in the form

$$\Psi_j(x,t) = e^{-i\omega t} \begin{pmatrix} A_j(x) \\ iB_j(x) \end{pmatrix}. \tag{10}$$

Then, from Eqs. (2) and (10) we have

$$\frac{dA_j}{dx} + (m+\omega)B_j - g_j^2(A_j^2 - B_j^2)B_j = 0, \tag{11}$$

$$\frac{dB_j}{dx} + (m-\omega)A_j - g_j^2(A_j^2 - B_j^2)A_j = 0. \tag{12}$$

The vertex boundary conditions for the functions A_j and B_j can be written as

$$\begin{aligned} \alpha_1 A_1|_{x=0} &= \alpha_2 A_2|_{x=0} + \alpha_3 A_3|_{x=0}, \\ \frac{1}{\alpha_1} B_1|_{x=0} &= \frac{1}{\alpha_2} B_2|_{x=0} = \frac{1}{\alpha_3} B_3|_{x=0}. \end{aligned} \tag{13}$$



FIG. 2. (Color online) Standing wave soliton solutions Eq.(2) described by Eqs. (14) and(15) for $g_1 = 1,\ g_2 = g_3 = \sqrt{2}$.

A prototypical static solution of the system (11), (12) obeying the asymptotic conditions (7) can be written as[26]

$$\begin{aligned} A_j(x) = &\sqrt{\frac{(m+\omega)\cosh^2(\beta x)}{m+\omega\cosh(2\beta x)}} \cdot \\ &\times \sqrt{\frac{2\beta^2}{g_j^2(m+\omega\cosh(2\beta x))}}, \end{aligned} \tag{14}$$

$$B_j(x) = \sqrt{\frac{(m-\omega)\sinh^2(\beta x)}{m+\omega\cosh(2\beta x)}}\cdot \times\sqrt{\frac{2\beta^2}{g_j^2(m+\omega\cosh(2\beta x))}}, \tag{15}$$

where $\beta = \sqrt{m^2-\omega^2}$. In order for these solutions of Eqs. (11) and (12) to solve the problem on the metric graph, they need to also satisfy the vertex boundary conditions (13). This can be achieved if the constants α_j and coupling parameters g_j fulfill the following conditions:

$$\frac{\alpha_1}{|g_1|} = \frac{\alpha_2}{|g_2|} + \frac{\alpha_3}{|g_3|}. \tag{16}$$

which stems from the first of Eqs. (9) and

$$\frac{\alpha_{2,3}}{\alpha_1} = \frac{|g_1|}{|g_{2,3}|}, \tag{17}$$

in accordance with the second one of conditions (9). The combination of the two leads to the "sum rule":

$$\frac{1}{g_1^2} = \frac{1}{g_2^2} + \frac{1}{g_3^2}. \tag{18}$$

It is important to note that this sum rule is derived by assuming that incoming wave comes from the first bond, b_1, while if it comes from the bond b_2 (or b_3) one should replace g_1 in Eq. (18) with g_2 (or g_3). In Fig. 2 plots of the soliton solutions of NLDE on metric star graph corresponding to Eqs. (14) and (15), satisfying the vertex conditions are given.

We note that soliton solutions given by Eqs.(14) and (15) describe the standing waves in metric graphs. The traveling wave (soliton) solutions of NLDE can be obtained by considering the case of the moving frame and taking into account the Lorentz invariance[26]. Introducing the Lorentz transformation between the frames moving with relative velocity, v in the form[26]

$$v = \tanh\eta,\ \gamma = \frac{1}{\sqrt{1-v^2}} = \cosh\eta,\ \sinh\eta = \frac{v}{\sqrt{1-v^2}},$$

one can obtain traveling wave (soliton) solutions of the nonlinear Dirac equation in the moving frame determined by the constraints (16). It is important to note that this transformation does not affect the scaling dependence on the coupling parameters g_j. For that reason, we expect the form of Eqs. (16) to still be valid in the case of traveling/moving wave solutions.

To analyze the dynamics of the Dirac solitons on metric graph, we solve numerically Eq. (2) for the vertex boundary conditions given by Eq. (9) by considering the moving frame. We do this for the case when the constraint given by Eq. (18) is fulfilled, as well as for the case of arbitrary g_j. Fig. 3(a) presents the plots of the solution of Eqs. (2) and (9) obtained numerically for the values of g_j obeying the constraint (18). To obtain numerical traveling wave solutions, the initial conditions are imposed as soliton solutions for the moving frame at $t = 0$. One can observe the absence of the vertex reflection in this case. Fig. 3(b) presents soliton solutions of nonlinear Dirac equation for the boundary conditions (9) obtained numerically for those values of g_j which do not fulfill Eq. (18). Fig. 4(a) shows the conservation of the energy and reflectionless transmission through the Y-junction during the propagation of Dirac soliton in graph. The emergence of the vertex reflection when the constraints (18) are not fulfilled, can also be seen from the Fig. 4(b) where the vertex reflection coefficient is plotted as a function of g_3 for fixed values of g_1 and g_2. This systematic analysis clearly illustrates the necessity of the symmetric scenario of $g_3 = 1/\sqrt{2}$ for reflection to be absent.



FIG. 3. (Color online) Numerical solution of Eq. (2) for boundary conditions (9) for the case (a) when constraints (18) are fulfilled ($g_1 = 1,\ g_2 = g_3 = \sqrt{2}$), (b) when these constraints are not fulfilled ($g_1 = 1,\ g_2 = \sqrt{2}; g_3 = 3.5$).



FIG. 4. (Color online) Numerical solution of Eq.(2) for boundary conditions (9). (a) Total energy and time-dependence of the energy for each bond with $g_1 = 1,\ g_2 = g_3 = \sqrt{2}$. (b) Reflection coefficient as a function of g_3 for $g_1 = 1,\ g_2 = \sqrt{2}$.

IV. EXTENSION FOR OTHER GRAPHS

The above treatment of the NLDE on the metric star graph can be extended to the tree graph presented in Fig. 5. It consists of three subgraphs $b_1, (b_{1i}), (b_{1ij})$, where i, j run over the given bonds of a subgraph. On each bond b_1, b_{1i}, b_{1ij} we posit that the nonlinear Dirac equation is satisfied as given by Eq. (2). The vertex boundary conditions can be written similarly to those in Eq. (13). Soliton solutions have a similar form to those in Eqs. (14)-(15) where x is replaced by $x + s_b$, with $s_1 = s_{1i} = l$, $s_{1ij} = l + L_i$, with l being the initial position of the center of mass of soliton.



FIG. 5. Tree graph. $x_1 \in (-\infty, 0], x_{1i} \in [0, L_i]$, and $x_{1ij} \in [0, +\infty)$ with $i = 1, 2; j = 1, 2, 3$.

The sum rules in this case generalize according to:

$$\frac{1}{g_1^2} = \sum_{i=1}^{2} \frac{1}{g_{1i}^2}, \; \frac{1}{g_{1i}^2} = \sum_{j=1}^{3} \frac{1}{g_{1ij}^2}, \; i = 1, 2.$$



FIG. 6. Triangle graph. $x_1 \in (-\infty, 0], x_i \in [0, L_i]$ with $i = 2, 3, 4$, and $x_j \in [0, +\infty)$ with $j = 5, 6$.

Similarly to that for star graph, one can obtain soliton solutions of nonlinear Dirac equation on the tree graph for the moving frame.

Another graph for which a soliton solution of the nonlinear Dirac equation can be obtained, is presented in Fig. 6. It has the form of a triangle whose vertices are connected to outgoing semi-infinite leads. The vertex boundary conditions for Eqs. (11) and (12) which follow from the conservation of current and energy can be written as

$$\begin{aligned}
\alpha_1 \phi_1|_{x=0} &= \alpha_2 \phi_2|_{x=0} + \alpha_3 \phi_3|_{x=0}, \\
\frac{1}{\alpha_1} \chi_1|_{x=0} &= \frac{1}{\alpha_2} \chi_2|_{x=0} = \frac{1}{\alpha_3} \chi_3|_{x=0}, \\
\alpha_2 \phi_2|_{x=L_2} &= \alpha_4 \phi_4|_{x=0} + \alpha_5 \phi_5|_{x=0}, \\
\frac{1}{\alpha_2} \chi_2|_{x=L_2} &= \frac{1}{\alpha_4} \chi_4|_{x=0} = \frac{1}{\alpha_5} \chi_5|_{x=0}, \\
\alpha_3 \phi_3|_{x=L_3} + \alpha_4 \phi_4|_{x=L_4} &= \alpha_6 \phi_6|_{x=0}, \\
\frac{1}{\alpha_3} \chi_3|_{x=L_3} &= \frac{1}{\alpha_4} \chi_4|_{x=L_4} = \frac{1}{\alpha_6} \chi_6|_{x=0}.
\end{aligned}$$



FIG. 7. (Color online) Numerical solution of Eq.(2) for the case (a) when constraints (21)–(23) are fulfilled ($g_1 = 1,\ g_2 = g_3 = \sqrt{2},\ g_4 = g_5 = 2,\ g_6 = 2/\sqrt{3}$), (b) when these constraints are not fulfilled ($g_1 = g_3 = g_5 = 1,\ g_2 = g_4 = g_6 = 2$).

In short, the conditions matching the flux of particle current and energy must be applied to each node connecting multiple vertices of the graph. The soliton solution obeying these boundary conditions can be written as

$$A_b(x) = \frac{1}{|g_b|}\cosh(\beta(x+s_b))\sqrt{\frac{m+\omega}{m+\omega\cosh(2\beta(x+s_b))}} \times\sqrt{\frac{2\beta^2}{m+\omega\cosh(2\beta(x+s_b))}}, \tag{19}$$

$$B_b(x) = \frac{1}{|g_b|}|\sinh(\beta(x+s_b))|\sqrt{\frac{m-\omega}{m+\omega\cosh(2\beta(x+s_b))}} \times\sqrt{\frac{2\beta^2}{m+\omega\cosh(2\beta(x+s_b))}}, \tag{20}$$

where b is considered to be an index running from 1 to 6, $s_1 = s_2 = s_3 = l$, $s_4 = l + L_2$, $s_5 = l + L_2$, $s_6 = l + L_3$, $L_3 = L_2 + L_4$ (this condition is necessary for the waves traveling along b_3 and those split along b_2 and b_4 to arrive at the

vertex between b_3 and b_4 concurrently) and coefficients g_j fulfill the following constraints:

$$\frac{1}{g_1^2} = \frac{1}{g_2^2} + \frac{1}{g_3^2}, \tag{21}$$

$$\frac{1}{g_2^2} = \frac{1}{g_4^2} + \frac{1}{g_5^2}, \tag{22}$$

$$\frac{1}{g_3^2} + \frac{1}{g_4^2} = \frac{1}{g_6^2}. \tag{23}$$

Again, the traveling wave (soliton) solutions can be obtained in case of the moving frame using the Lorentz transformations. This approach for designing soliton solutions of the nonlinear Dirac equation can be applied to other graph topologies with multiple junctions, provided a graph consists of finite parts connected to outgoing semi-infinite bonds.

In Fig. 7 the solution of NLDE on each bond of this triangle graph is plotted for the cases when the sum rule given by Eqs. (21)-(23) is fulfilled (a) and is broken (b). Absence of the vertex reflection in Fig. 7(a) is clearly seen, while in Fig. 7(b) the transmission of Dirac solitons through the graph vertices is clearly impeded by the mismatch at the VBC. Thus, reflection events are clearly discernible in bonds such as 1, 3 and 4.

V. CONCLUSION

In this paper we studied dynamics of Dirac solitons in networks by considering the case of metric graphs. We obtained soliton solutions of the nonlinear Dirac equation on some of the simplest metric graphs such as the star, tree and triangle graphs. Constraints allowing such exact solutions are derived in the form of sum rules for bond nonlinearity coefficients. The boundary conditions at the branching points (vertices) are derived from the fundamental conservation laws. It is shown via direct numerical simulations that the obtained solutions provide for reflectionless transmission of solitons at the graph vertex. In the case where the relevant conditions are violated, nontrivial reflections ensue.

Naturally, the above study can be extended for other simple topologies, such as a graph with one or multiple loops (e.g. the dumbbell graph), or other combinations of star, tree and loop graphs. It will be extremely interesting if these ideas induce experimental interest in systems such as suitably tailored discrete optical waveguides[50,51] or in atomic settings, in the same way as they have for instance for the propagation of traveling waves through networks of granular crystals[52].

ACKNOWLEDGEMENT

The work of KKS is partially supported by a grant of the Committee for Coordination of Science and Technology of Uzbekistan (Ref. Nr F-2-003).

VI. APPENDIX A

Here we will show that the linear vertex boundary conditions given by Eqs. (9) lead to the ones given by (6)-(8). Consider the following (linear) relations given at the vertex of the a metric star graph

$$\alpha_1\phi_1|_{x=0} = \alpha_2\phi_2|_{x=0} + \alpha_3\phi_3|_{x=0}, \tag{24}$$

$$\frac{1}{\alpha_1}\chi_1|_{x=0} = \frac{1}{\alpha_2}\chi_2|_{x=0} = \frac{1}{\alpha_3}\chi_3|_{x=0}, \tag{25}$$

where α_1, α_2, α_3 are real constants.

Multiplying both sides of Eq. (24) by $\frac{1}{\alpha_1}\chi_1^*|_{x=0}$, from Eq. (25) we have

$$\phi_1\chi_1^*|_{x=0} = \phi_2\chi_2^*|_{x=0} + \phi_3\chi_3^*|_{x=0}. \tag{26}$$

From Eq. (26) we get

$$\mathrm{Re}[\phi_1\chi_1^*]|_{x=0} = \mathrm{Re}[\phi_2\chi_2^*]|_{x=0} + \mathrm{Re}[\phi_3\chi_3^*]|_{x=0}, \tag{27}$$

which is nothing but the vertex boundary condition given by Eq. (6). Taking the time derivative from Eq. (25) and multiplying both sides of (24) by $\frac{1}{\alpha_1}\partial_t\chi_1^*|_{x=0}$, we obtain

$$\phi_1\partial_t\chi_1^*|_{x=0} = \phi_2\partial_t\chi_2^*|_{x=0} + \phi_3\partial_t\chi_3^*|_{x=0}. \tag{28}$$

Taking the time-derivative from Eq. (24) and multiplying the complex conjugate of the obtained expression by $\frac{1}{\alpha_1}\chi_1|_{x=0}$ we get

$$\chi_1\partial_t\phi_1^*|_{x=0} = \chi_2\partial_t\phi_2^*|_{x=0} + \chi_3\partial_t\phi_3^*|_{x=0}. \tag{29}$$

Adding Eqs. (28) and (29) leads to

$$\begin{aligned}[\phi_1\partial_t\chi_1^* + \chi_1\partial_t\phi_1^*]|_{x=0} = {} & [\phi_2\partial_t\chi_2^* + \chi_2\partial_t\phi_2^*]|_{x=0} \\ & + [\phi_3\partial_t\chi_3^* + \chi_3\partial_t\phi_3^*]|_{x=0}.\end{aligned} \tag{30}$$

The last equation yields the vertex boundary condition given by Eq. (8).

[1] Z. Sobirov, D. Matrasulov, K. Sabirov, S. Sawada, and K. Nakamura, Phys. Rev. E **81**, 066602 (2010).
[2] K. Nakamura, Z. A. Sobirov, D. U. Matrasulov, and S. Sawada, Phys. Rev. E **84**, 026609 (2011).
[3] R. Adami, C. Cacciapuoti, D. Finco, and D. Noja, Rev. Math. Phys. **23**, 409 (2011).
[4] R. Adami, C. Cacciapuoti, D. Finco, and D. Noja, Europhys. Lett. **100**, 10003 (2012).
[5] R. Adami, D. Noja, and C. Ortoleva, J. Math. Phys. **54**, 013501 (2013).
[6] K.K. Sabirov, Z.A. Sobirov, D. Babajanov, and D.U. Matrasulov, Phys. Lett. A, **377**, 860 (2013).
[7] D. Noja, Philos. Trans. R. Soc. A **372**, 20130002 (2014).
[8] G. H. Susanto and S.A. van Gils, Phys. Lett. A 338, 239 (2005).
[9] J.-G Caputo , D. Dutykh, Phys. Rev. E **90**, 022912 (2014).
[10] H. Uecker, D. Grieser, Z. Sobirov, D. Babajanov, and D. Matrasulov, Phys. Rev. E **91**, 023209 (2015).
[11] Z. Sobirov, D. Babajanov, D. Matrasulov, K. Nakamura, and H. Uecker, EPL **115** , 50002 (2016).
[12] W. Thirring, Ann. Phys. **3**, 91 (1958).
[13] D. J. Gross, A. Neveu, Phys. Rev. D **10**, 3235 (1974).
[14] W.H. Steeb, W. Oevel, and W. Strampp, J. Math. Phys. **25**, 2331 (1984).
[15] W.I. Fushchych and R.Z. Zhdanov, J. Math. Phys. **32**, 3488 (1991).
[16] F.M. Toyama, Y. Hosono, B. Ilyas and Y. Nogami, J. Phys. A: Math. Gen. **27**, 3139 (1994).
[17] W.K. Ng, Int. J. Mod. Phys. A, **24**, 3476 (2009).
[18] W.K. Ng and R.R. Parwani, SIGMA **5**, 023 (2009).
[19] L. H. Haddad and L. D. Carr, Physica D **238**, 1413 (2009).
[20] L. H. Haddad and L. D. Carr, Europhys. Lett. **94** 56002 (2011).
[21] L.H. Haddad, C.M. Weaver and L.D Carr, New J. Phys. **17**, 063033 (2015).
[22] L.H. Haddad, C.M. Weaver and L.D Carr, New J. Phys. **17**, 063034 (2015)
[23] L.H. Haddad, C.M. Weaver and L.D Carr, New J. Phys. **17**, 113011 (2015)
[24] L.H. Haddad, K. M. O'Hara, L.D Carr, Phys. Rev. A **91**, 043609 (2015).
[25] F.Cooper, A. Khare, B. Mihaila and A. Saxena, Phys. Rev. E **82**, 036604 (2010).
[26] F. G. Mertens, N. R. Quintero, F. Cooper, A. Khare, and A. Saxena, Phys. Rev. E **86**, 046602 (2012).
[27] D.E. Pelinovsky, A.Stefanov, J. Math.Phys.,**53**, 073705 (2012).
[28] D.E. Pelinovsky, Y. Shimabukuro, Lett. Mat. Phys.,**104**, 21 (2014).
[29] S. Shao, F. G. Mertens, N. R. Quintero, F. Cooper, A. Khare, and A. Saxena, Phys. Rev. E **90**, 032915 (2014).

[30] J. Cuevas-Maraver, P.G Kevrekidis and A. Saxena, J. Phys. A: Math. Gen. **48**, 055204 (2015).
[31] F. G. Mertens, F. Cooper, N.R. Quintero, S. Shao, A. Khare, and A. Saxena, J. Phys. A: Math. Gen. **49**, 065402 (2016).
[32] J. Cuevas-Maraver, P.G. Kevrekidis and A. Saxena, F. Cooper, A. Khare, A. Comech, C.M.Bender, IEEE J. Selected Topics Quant. Electr.,**22**, 5000109 (2016).
[33] J. Cuevas-Maraver, P.G Kevrekidis and A. Saxena, A.Comech, R.Lan, Phys. Rev. Lett. **116**, 214101 (2016).
[34] A. Contreras, D.E. Pelinovsky, Y. Shimabukuro, Commun. Part. Diff. Eq.,**41**, 227 (2016).
[35] A. Komech, A. Komech, SIAM J. Math. Anal., **42**, 2944 (2010).
[36] G. Berkolaiko, A. Comech, Math. Mod. of Natural Phenomena, **7**, 13 (2012).
[37] A. Comech, M. Guan and S. Gustafson, Ann. Henri Poincare (Analyse non lineaire), **31** 639 (2014).
[38] G. Berkolaiko, A. Comech, A. Sukhtayev, Nonlinearity, **28** 577 (2015).
[39] J. Cuevas-Maraver, P.G. Kevrekidis, A. Saxena, A. Comech, R. Lan, Phys. Rev. Lett. **116** 214101 (2016).
[40] Y Zhang, Nonlinear Analysis, **118** 82 (2015).
[41] Ch-W. Lee, P. Kurzynski, H. Nha, Phys. Rev. A **92** 052336 (2015).
[42] T.X. Tran, X.N. Nguyen, F. Biancalana, Phys. Rev. A **91**, 023814 (2008).
[43] A.N. Andriotis, M. Menon, Appl.Phys.Lett., **92** 042115 (2008).
[44] Y.P. Chen, Y.E. Xie, L. Z. Sun and J. Zhong, Appl. Phys. Lett., **93** 092104 (2008).
[45] J. G. Xu, L. Wang and M. Q. Weng, J. Appl. Phys., **114** 153701 (2013).
[46] A.G. Kvashnin, D.G. Kvashnin, O.P. Kvashnina and L.A. Chernozatonskii, Nanotechnology, **26** 385705 (2015).
[47] T.X. Tran, S. Longhi, F. Biancalana, Ann. Phys. **340** 179 (2014).
[48] D.N. Christodoulides and E.D. Eugenieva Phys. Rev. Lett. **87**, 233901 (2001)
[49] J. Bolte and J. Harrison, J. Phys. A: Math. Gen. **36** 2747 (2003).
[50] M. Conforti, C. De Angelis, T. R. Akylas, and A.B. Aceves Phys. Rev. A **85** 063836 (2012).
[51] A. Marini, S. Longhi, and F. Biancalana Phys. Rev. Lett. **113** 150401 (2014).
[52] A. Leonard, L. Ponson, C. Daraio, J. Mech. Phys. Solids **73**, 103 (2014).